




Lake Mitchell 2021 Aquatic Vegetation, Water Quality, and 2022 Management Recommendations Report



January, 2022

Lake Mitchell 2021 Aquatic Vegetation, Water Quality, and 2022 Management Recommendations Report (2009-2021)



**© Restorative Lake Sciences
18406 West Spring Lake Road
Spring Lake, Michigan 49456
Phone 616.843.5636**

Website: <http://www.restorativelakesciences.com>

Table of Contents

Section 1: Lake Mitchell Summary (2021)	4
Section 2: Lake Mitchell Water Quality Data (2009-2021)	5
Section 3: Lake Mitchell Aquatic Vegetation Data (2021)	13
Section 4: Management Recommendations for 2022	21

Lake Mitchell 2021 Aquatic Vegetation, Water Quality, and 2022 Management Recommendations

The overall condition of Lake Mitchell in 2021 was good with favorable water clarity, reduced total phosphorus concentrations, and reduced EWM growth. The water clarity in 2021 averaged around 9.5 feet which is favorable. The reduced EWM growth is a result of rigorous whole-lake surveys, intermittent site visits, and rigorous applications of systemic herbicides for the target plants. In 2021, RLS utilized the new systemic herbicide ProcellaCOR® along with diquat with excellent success and that is recommended for 2022. RLS is always present to oversee these herbicide treatments to assure that exact locations are addressed. Over the past decade, the LMIB has saved much needed funds due to these efforts. The EWM in Big Cove sustained reduced growth in 2021 after rigorous 2020 ProcellaCOR® treatments.

Protection of the 26 native aquatic plant species is paramount for the health of the lake fishery and these plants should not be managed unless they are a nuisance to lakefront property owners and possess navigational and recreational hazards (i.e., lily pads or nuisance pondweeds in the coves).

Invasive species such as Eurasian Watermilfoil (EWM) are able to grow in moderate nutrient waters and thus are a challenge to the Lake Mitchell ecosystem. In 2021, approximately 64.6 acres of EWM were treated throughout the entire lake. RLS has recommended alternating use of different systemic herbicides to reduce the probability of herbicide tolerance which reduces efficacy. The Purple Loosestrife beetle stocking is recommended in 2022 to increase control of the plant or spot-treatments with an aquatic herbicide for emergents. A thorough section on management recommendations for 2022 is offered at the end of this report.

Lake Mitchell Water Quality Data (2009-2021)

Water Quality Parameters Measured

There are numerous water quality parameters one can measure on an inland lake, but several are the most critical indicators of lake health. These parameters include water temperature (measured in °C), dissolved oxygen (measured in mg/L), pH (measured in standard units-SU), conductivity (measured in micro-Siemens per centimeter- $\mu\text{S}/\text{cm}$), total alkalinity or hardness (measured in mg of calcium carbonate per liter-mg CaCO_3/L), total dissolved solids (mg/L), secchi transparency (feet), total phosphorus chlorophyll-*a* (in $\mu\text{g}/\text{L}$), and algal species composition. Water quality was measured in the deepest basins of Lake Mitchell on September 21, 2021 (Figure 1). Trend data was calculated using mean values for each parameter for each season. Lake Mitchell would be considered eutrophic (relatively productive) since it does contain ample phosphorus, nitrogen, and aquatic vegetation growth but also has good water clarity and moderate algal growth. General water quality classification criteria are defined in Table 1. 2021 water quality data for Lake Mitchell and its tributaries are shown below in Tables 2-4.

Figure 1. Water quality sampling locations for Lake Mitchell and its tributaries




Table 1. Lake trophic classification (MDNR).

<i>Lake Trophic Status</i>	<i>Total Phosphorus ($\mu\text{g L}^{-1}$)</i>	<i>Chlorophyll-a ($\mu\text{g L}^{-1}$)</i>	<i>Secchi Transparency (feet)</i>
Oligotrophic	< 10.0	< 2.2	> 15.0
Mesotrophic	10.0 – 20.0	2.2 – 6.0	7.5 – 15.0
Eutrophic	> 20.0	> 6.0	< 7.5

Table 2. Lake Mitchell water quality parameter data collected over the north deep basin on September 21, 2021.

<i>Depth ft.</i>	<i>Water Temp $^{\circ}\text{C}$</i>	<i>DO mg L^{-1}</i>	<i>pH S.U.</i>	<i>Cond. $\mu\text{S cm}^{-1}$</i>	<i>Turb. NTU</i>	<i>ORP mV</i>	<i>Total Dissolved Solids mg L^{-1}</i>	<i>Total Alk. $\text{mg L}^{-1} \text{CaCO}_3$</i>	<i>Total Phos. mg L^{-1}</i>	<i>TKN mg L^{-1}</i>
0	20.3	8.8	8.0	152	0.6	135.6	98	46	0.027	0.5
10	20.3	8.5	8.0	152	0.6	122.6	98	46	0.028	0.6
19	20.3	8.5	7.9	153	1.8	101.5	98	47	0.027	<0.5

Table 3. Lake Mitchell water quality parameter data collected over the south deep basin on September 21, 2021.

<i>Depth ft.</i>	<i>Water Temp $^{\circ}\text{C}$</i>	<i>DO mg L^{-1}</i>	<i>pH S.U.</i>	<i>Cond. $\mu\text{S cm}^{-1}$</i>	<i>Turb. NTU</i>	<i>ORP mV</i>	<i>Total Dissolved Solids mg L^{-1}</i>	<i>Total Alk. $\text{mg L}^{-1} \text{CaCO}_3$</i>	<i>Total Phos. mg L^{-1}</i>	<i>TKN mg L^{-1}</i>
0	19.7	8.9	8.0	153	0.6	139.8	98	46	0.024	0.6
10	19.7	8.7	8.0	153	0.9	105.1	98	46	0.032	0.6
20	19.7	8.7	8.0	153	2.1	100.4	98	46	0.027	0.5


Table 4. Lake Mitchell Tributary water quality parameter data collected on September 21, 2021.

<i>Tributary</i>	<i>Water Temp °C</i>	<i>DO mg L⁻¹</i>	<i>pH S.U.</i>	<i>Cond. µS cm⁻¹</i>	<i>TDS mg L⁻¹</i>	<i>Total Phos. mg L⁻¹</i>	<i>TKN mg L⁻¹</i>
Mitchell	18.2	8.7	7.5	182	117	0.035	0.8
Brandy	17.3	6.7	7.1	104	67	0.160	1.4
Gyttja	19.4	8.2	7.2	177	113	0.029	0.9

Water Clarity (Transparency) Data


Elevated Secchi transparency readings allow for more aquatic plant and algae growth. The transparency in Lake Mitchell during the 2021 September sampling event averaged around 7.5 feet which was higher than observed in 2021. Earlier season measurements ranged from 10-13 feet with an overall mean of 9.5 feet for the season. Secchi transparency is variable and depends on the number of suspended particles in the water (often due to windy conditions of lake water mixing) and the amount of sunlight present at the time of measurement. Other parameters such as turbidity (measured in NTU's) and total dissolved solids (measured in mg/L) are correlated with water clarity and show an increase as clarity decreases. The turbidity and total dissolved solids in Lake Mitchell were quite low in 2021 at ≤2.1 NTU's and ≤98 mg/L, respectively. The graph below shows the trend in mean Secchi transparency over time for Lake Mitchell.

Trend in Lake Mitchell Mean Secchi Transparency



Total Phosphorus


Total phosphorus (TP) is a measure of the amount of phosphorus (P) present in the water column. Phosphorus is the primary nutrient necessary for abundant algae and aquatic plant growth. TP concentrations are usually higher at increased depths due to higher release rates of P from lake sediments under low oxygen (anoxic) conditions. Phosphorus may also be released from sediments as pH increases. Fortunately, even though the TP levels in Lake Mitchell are moderate, the dissolved oxygen levels are good enough at the bottom to not cause release of phosphorus from the bottom. TP concentrations during the 2021 sampling event ranged from 0.024-0.032 mg L⁻¹, with the highest concentration at the middle of the south basin (below figure).



Total Alkalinity

Lakes with high alkalinity ($> 150 \text{ mg L}^{-1}$ of CaCO_3) are able to tolerate larger acid inputs with less change in water column pH. Many Michigan lakes contain high concentrations of CaCO_3 and are categorized as having “hard” water. Total alkalinity may change on a daily basis due to the re-suspension of sedimentary deposits in the water and respond to seasonal changes due to the cyclic turnover of the lake water. The alkalinity of Lake Mitchell is quite low and is indicative of a “soft water” aquatic ecosystem. The total alkalinity during the sampling event in 2021 ranged from 46-47 mg L^{-1} of CaCO_3 which is similar to recent years (below figure).


Trend in Lake Mitchell Mean Total Alkalinity



pH


Most Michigan lakes have pH values that range from 6.5 to 9.5. Acidic lakes (pH < 7) are rare in Michigan and are most sensitive to inputs of acidic substances due to a low acid neutralizing capacity (ANC). Lake Mitchell is considered “neutral” on the pH scale. The pH of Lake Mitchell in 2021 was similar to previous years and was 8.0 S.U. (below figure).

Trend in Lake Mitchell Mean pH



Conductivity


Conductivity is a measure of the number of mineral ions present in the water, especially those of salts and other dissolved inorganic substances. Conductivity generally increases as the amount of dissolved minerals and salts in a lake increases, and also increases as water temperature increases. The conductivity values for Lake Mitchell are moderately low for a large, shallow inland lake and were all recorded at 152-153 $\mu\text{S}/\text{cm}$ during the 2021 sampling event which is lower than previous years (below figure). Severe water quality impairments do not occur until values exceed 800 $\mu\text{S}/\text{cm}$ and are toxic to aquatic life around 1,000 $\mu\text{S}/\text{cm}$. Conductivity may be increasing due to more road salt applications during recent harsh winters.



Chlorophyll-*a* and Algal Species Composition

Chlorophyll-*a* is a measure of the amount of green plant pigment present in the water, often in the form of planktonic algae. High chlorophyll-*a* concentrations are indicative of nutrient-enriched lakes. Chlorophyll-*a* concentrations greater than 6 $\mu\text{g L}^{-1}$ are found in eutrophic or nutrient-enriched aquatic systems, whereas chlorophyll-*a* concentrations less than 2.2 $\mu\text{g/L}$ are found in nutrient-poor or oligotrophic lakes. The mean chlorophyll-*a* concentrations on September 21, 2021 in Lake Mitchell was 7.5 $\mu\text{g/L}$ which was elevated for an inland Michigan lake and higher than in recent years (below figure).

The algal genera were determined from composite water samples collected over the deep basins of Lake Mitchell in 2021 were analyzed with a compound bright field microscope. The genera present included the Chlorophyta: *Chlorella* sp., *Haematococcus* sp., *Scenedesmus* sp., *Rhizoclonium* sp., *Mougeotia* sp., *Cladophora* sp., *Spirogyra* sp., *Staurastrum* sp., *Pediastrum* sp., and *Chloromonas* sp. The Cyanophyta (blue-green algae): *Microcystis* sp.; The Bascillariophyta (diatoms): *Synedra* sp., *Cymbella* sp., *Navicula* sp., and *Fragilaria* sp. The aforementioned species indicate a diverse algal flora and represent a good diversity of alga with an abundance of diatoms that are indicative of good water quality.



Aquatic Vegetation Data (2021)

Status of Native Aquatic Vegetation in Lake Mitchell

The native aquatic vegetation present in Lake Mitchell is essential for the overall health of the lake and the support of the lake fishery. The September 21, 2021 all species inventory survey determined that there were a total of 26 native aquatic plant species in Lake Mitchell. These include 17 submersed species, 4 floating-leaved species, and 5 emergent species which is similar to recent years and means that the lake is maintaining its biodiversity. This indicates a very high biodiversity of aquatic vegetation in Lake Mitchell and is likely a significant reason for the great fishery in the lake. The overall % cover of the lake by native aquatic plants is low relative to the lake size and thus these plants should be protected and not treated unless they become a nuisance in shallow coves or the Torenta Canal. In these cases, RLS may recommend the use of mechanical harvesting in some areas of Big Cove and/or Little Cove along with the Torenta Canal. A list of all native aquatic plants and their relative abundance can be found in Table 5 below.

The most common aquatic plants found during the 2021 surveys included: 1) White-stem Pondweed which has long, green leaves and a prominent seed head that often breaks the lake surface; 2) Chara, which is a macro alga and is brittle in texture with a strong sulfur odor, and 3) Fern-leaf Pondweed, which grows close the lake bottom and has fern-like leaves that may have a green or brown hue.

During the whole-lake scan, an aquatic vegetation biovolume map (Figure 4) was developed which shows the areas where aquatic vegetation is absent (blue color), sparse (green color), or high-growing (red color). The red colors usually represent milfoil growth in Lake Mitchell which has declined over the past few years.

Table 5. Native aquatic plants found in Lake Mitchell on September 21, 2021.

<i>Aquatic Plant Species Name</i>	<i>Aquatic Plant Common Name</i>	<i>Aquatic Plant Growth Form</i>	<i>% Coverage of Lake (2021)</i>
<i>Chara vulgaris</i> (macroalga)	Muskgrass	Submersed; Rooted	21.9
<i>Potamogeton pectinatus</i>	Sago Pondweed	Submersed; Rooted	2.1
<i>Potamogeton robbinsii</i>	Fern-leaf Pondweed	Submersed; Rooted	13.0
<i>Potamogeton gramineus</i>	Variable-leaf Pondweed	Submersed; Rooted	6.0
<i>Potamogeton praelongus</i>	White-stem Pondweed	Submersed; Rooted	42.1
<i>Potamogeton richardsonii</i>	Clasping-leaf Pondweed	Submersed; Rooted	7.0
<i>Potamogeton illinoensis</i>	Illinois Pondweed	Submersed; Rooted	8.0
<i>Potamogeton amplifolius</i>	Large-leaf Pondweed	Submersed; Rooted	9.2
<i>Myriophyllum sibiricum</i>	Northern Watermilfoil	Submersed; Rooted	1.8
<i>Ceratophyllum demersum</i>	Coontail	Submersed; Non-rooted	0.8
<i>Elodea canadensis</i>	Common Waterweed	Submersed; Rooted	4.5
<i>Utricularia vulgaris</i>	Common Bladderwort	Submersed; Non-rooted	0.9
<i>Utricularia minor</i>	Mini Bladderwort	Submersed; Non-rooted	0.1
<i>Najas guadalupensis</i>	Southern Naiad	Submersed; Rooted	12.0
<i>Najas flexilis</i>	Slender Naiad	Submersed; Rooted	8.0
<i>Myriophyllum tenellum</i>	Leafless Watermilfoil	Submersed; Rooted	21.5
<i>Potamogeton pusillus</i>	Small-leaf Pondweed	Submersed; Rooted	2.0
<i>Megalodonta beckii</i>	Water Marigold	Submersed; Rooted	2.5
<i>Nymphaea odorata</i>	White Waterlily	Floating-leaved	8.0
<i>Nuphar variegata</i>	Yellow Waterlily	Floating-leaved	8.5
<i>Brasenia schreberi</i>	Watershield	Floating-leaved	9.1
<i>Lemna trisulca</i>	Star Duckweed	Floating-Leaved; Non-rooted	0.1
<i>Pontedaria cordata</i>	Pickerelweed	Emergent	9.2
<i>Typha latifolia</i>	Cattails	Emergent	8.1
<i>Schoenoplectus acutus</i>	Bulrushes	Emergent	15.0
<i>Decodon verticillatus</i>	Swamp Loosestrife	Emergent	11.6
<i>Eleocharis acicularis</i>	Spike rush	Emergent	7.2




Figure 4. Aquatic Vegetation Biovolume in Lake Mitchell (September 21, 2021).

Status of Invasive (Exotic) Aquatic Plant Species in Lake Mitchell

The amount of Eurasian Watermilfoil (Figure 5) present in Lake Mitchell varies each year and is dependent upon climatic conditions, especially runoff-associated nutrients. A whole-lake survey of the main lake was conducted on May 25, 2021 and revealed that approximately 7.0 acres of milfoil were found throughout the entire lake. However, by June 1, 2021 that amount increased to 35.0 acres and was treated on June 3, 2021. Significant growth can occur when there is a large increase in air and water temperatures and high sunlight periods that allow for high sunlight penetration such as in spring, 2021. An additional survey on August 4, 2021 determined an additional 20 acres of EWM was present but by the August 19, 2021 treatment that increased by 4.6 acres and all EWM was treated. Surveys in between treatments allowed for determination of small areas of EWM growth that were prompt treated along with the coves. In 2021, Torenta Canal did not require treatment. Table 6 below shows the total acres of milfoil and nuisance weeds (Figure 6) found in each region of the lake that was treated on various dates. Also noted are the effective products and doses used.

The treatments were very successful with little viable milfoil remaining at the end of 2021. A spring 2022 survey is needed, however, to determine the 2021 treatment efficacy as EWM plants were senesced by late September 2021 but may re-germinate in the same locations in 2022. Treatment maps for each of these invasive species are shown in the maps below (Figures 7 and 8). Also noted are the effective products and doses used.



Figure 5. Eurasian Watermilfoil



Figure 6. Curly-leaf Pondweed

Table 6. Number of acres of nuisance aquatic vegetation managed inn various regions of Lake Mitchell (June-August 2021).

Area of Lake Treated	Date Treated	# Acres of EWM	# Acres of CLP or Nuisance Pondweeds	Products Used and Associated Doses
Franke South Cove	6-29	--	--	Phoslock Treatment (1,485 lbs.)
	7-12	--	3.6	Aquathol-K (3 gal/acre) + flumioxazin at 200 ppb
Main Lake	6-3	35.5	--	ProcellaCOR/diquat (4 PDU + 1 gal/acre)
	8-19	24.6		
Big Cove	8-19	4.6	--	ProcellaCOR/diquat (4 PDU + 1 gal/acre)
Little Cove	8-2	--	1.8	Aquathol-K (3.5 gals) + flumioxazin at 200 ppb
	8-19	3.0	--	ProcellaCOR/diquat (4 PDU + 1 gal/acre)
Franke North Cove	7-12	--	2.4	Aquathol-K (3 gal/acre) + flumioxazin at 200 ppb
	8-2	4.5	--	ProcellaCOR/diquat (18 PDU/4.5 gals)
Torenta Canal	--	--	--	--




Figure 7. Distribution of EWM in Lake Mitchell (May 25, 2021). A marked reduction in EWM in the main lake occurred relative to previous years due to intense treatment efforts and surveys.





Figure 8. Late-season Distribution of EWM and nuisance weeds in the main lake and coves (August 4, 2021)

Evaluation of Purple Loosestrife Beetles on Lake Mitchell Purple Loosestrife Reduction:

The beetle, *Galerucella* sp. is stocked each season around areas of Lake Mitchell infested with Purple Loosestrife. The goal has been to introduce enough beetles each season to create a sustainable population around the lake to naturally take over management of the invasive Purple Loosestrife. Beetle counts are performed on the plants each year to evaluate the number of beetles found along with damage of the inflorescences (flower portions of the plants). Based on the graph below, the beetle population is declining and did not have marked effects on the Purple Loosestrife in 2020-2021. More stocking is recommended for 2021-2 and beyond as budget allows or the LMIB may consider topical treatment of the plants with triclopyr.



Management Recommendations for 2022


Detailed aquatic vegetation surveys will be done in 2022 to determine locations of EWM, CLP, and any other nuisance invasive species or natives that may be a threat to the Lake Mitchell ecosystem. Along with the surveys, bottom scans will be conducted to determine changes in aquatic biovolume and distribution of aquatic vegetation. These surveys will occur during the summer or early fall depending on weather patterns which correspond with growth patterns. A post-treatment survey will also be scheduled, along with intermittent post-treatment surveys if small-scale treatments are conducted. RLS scientists will oversee all treatments as in previous years. RLS will notify the LMIB of the 2022 survey and treatment dates and update the LMIB on all management activities. Treatments may need to be rescheduled dependent upon wind and other weather conditions.

This year RLS is recommending that we treat large offshore areas with a combination of the systemic herbicide ProcellaCOR® along with the contact herbicide diquat, which resulted in sustained control on EWM in 2021. Diquat and/or flumioxazin will continue to be used in the cove areas for nuisance natives. An additional Phoslock treatment may be desired in Franke South and RLS supports this to reduce the nutrients that are resulting in enhanced weed and algae growth. The maximum allowed dosage of flumioxazin is now at 200 ppb per EGLE requirements. Maintaining EWM at existing low levels will be the top priority to keeping a healthy aquatic plant balance and continuing to maintain a low assessment for the lakefront owners in the special assessment district. The canal will be assessed for the need for a possible harvest and scheduled if necessary.

Water quality will continue to be monitored in the lake and tributaries. New water quality data from 2022 will be compared to historic data to continue evaluation of long-term trends. Lake Mitchell is a healthy lake with excellent aquatic plant diversity. Nutrients are at acceptable levels and there is a robust fishery indicated by the many fishing tournaments held on the lake.

Temporary algal blooms occur during hot windless periods or after intense rainfall events. RLS will continue to monitor the lake for any problematic algal blooms.

Lake Mitchell Improvement Board meetings will be attended by an RLS scientist as in previous years and RLS will develop a comprehensive annual report during the year that will be presented to the LMIB in the fall or winter of 2022. The graph below shows the results of the successful EWM reduction plan for Lake Mitchell which has resulted in substantial savings to the LMIB over the years.



Glossary of Scientific Terms used in this Report

- 1) Biodiversity- The relative abundance or amount of unique and different biological life forms found in a given aquatic ecosystem. A more diverse ecosystem will have many different life forms such as species.
- 2) CaCO_3 - The molecular acronym for calcium carbonate; also referred to as “marl” or mineral sediment content.
- 3) Eutrophic- Meaning “nutrient-rich” refers to a lake condition that consists of high nutrients in the water column, low water clarity, and an over-abundance of algae and aquatic plants.
- 4) Mesotrophic- Meaning “moderate nutrients” refers to a lake with a moderate quantity of nutrients that allows the lake to have some eutrophic qualities while still having some nutrient-poor characteristics
- 5) Oligotrophic- Meaning “low in nutrients or nutrient-poor” refers to a lake with minimal nutrients to allow for only scarce growth of aquatic plant and algae life. Also associated with very clear waters.
- 6) Sedimentary Deposits- refers to the type of lake bottom sediments that are present. In some lakes, gravel and sand are prevalent. In others, organic muck, peat, and silt are more common.